arXiv:1607.08423v1 [math.AP] 28 Jul 2016

The evolution to localized and front solutions in a non-Lipschitz reaction-diffusion Cauchy problem with trivial initial data

J. C. Meyer and D. J. Needham

July 29, 2016

Abstract

In this paper, we establish the existence of spatially inhomogeneous classical self-similar solutions to a non-Lipschitz semi-linear parabolic Cauchy problem with trivial initial data. Specifically we consider bounded solutions to an associated two-dimensional non-Lipschitz non-autonomous dynamical system, for which, we establish the existence of a two-parameter family of homoclinic connections on the origin, and a heteroclinic connection between two equilibrium points. Additionally, we obtain bounds and estimates on the rate of convergence of the homoclinic connections to the origin.

1 Introduction

In this paper, we study classical bounded solutions $u : \mathbb{R} \times [0,T] \to \mathbb{R}$ to the non-Lipschitz semi-linear parabolic Cauchy problem

$$u_t - u_{xx} = u|u|^{p-1} \quad \text{on } \mathbb{R} \times (0,T], \tag{1}$$

$$u = 0 \quad \text{on } \mathbb{R} \times \{0\}, \tag{2}$$

with $0 < p < 1$ and $T > 0$ (which we henceforth refer to as [CP]). The primary achievement of the paper is the establishment of the existence of a two-parameter family of localized spatially inhomogeneous solutions to [CP] for which $u(x,t) \to 0$ as $|x| \to \infty$ uniformly for $t \in [0,T]$; the secondary achievement of the paper is the establishment of front solutions to [CP], which approach $\pm(1-p)^{1/(1-p)}t$ as $|x| \to \pm\infty$ uniformly for $t \in [0,T]$. We note here that for $p \geq 1$ in (1), then the unique bounded classical solution with initial data (2) is the trivial solution, see for example [15, Theorem 4.5].

Qualitative properties of non-negative (non-positive) solutions to (1) when $0 < p < 1$, with non-negative (non-positive) initial data, and for which $u(x,t)$ is bounded as $|x| \to \infty$ uniformly for $t \in [0,T]$, have been determined in [1], [12], [20], [14] and [16]. However, we note that any non-negative (non-positive)

classical bounded solution to [CP] must be spatially homogeneous for $t \in [0,T]$, see for example [1, Corollary 2.6]. Thus, the solutions constructed in this paper are two signed on $\mathbb{R} \times [0,T]$. The authors are currently unaware of any studies of two signed solutions to (1)-(2) with $0 < p < 1$. Generic local results for spatial homogeneity of solutions to semi-linear parabolic Cauchy problems with homogeneous initial data depend upon uniqueness results, see for example, [16]. For results concerning the related problem of asymptotic homogeneity (in general, asymptotic symmetry) as $t \to \infty$ of non-negative (non-positive) global solutions to semi-linear parabolic Cauchy problems, we refer the reader to the survey article [22].

Non-negative (non-positive), spatially inhomogeneous solutions to (1) for $p > 1$ have been considered in [10], [26] [27], [21], [9], [11], [24], [8], [5] and [25] with the focus primarily on critical exponents for finite time blow-up of solutions, and conditions for the existence of global solutions (see the review articles [13] and [7]). Moreover, for $p > 1$, solutions to (1) with two signed initial data have been considered in [18] and [19], whilst boundary value problems have been studied in [3] and [4].

The paper is structured as follows; in Section 2 we introduce the self-similar solution structure for [CP], and hence, determine an ordinary differential equation related to (1); the remainder of the paper concerns the study of particular solutions to this ordinary differential equation, which is re-written as an equivalent two-dimensional non-autonomous dynamical system. Specifically, in Section 3 we establish the existence of a two-parameter family of homoclinic connections on the equilibrium $(0,0)$. Additionally, we determine bounds and estimates on the asymptotic approach of these solutions to $(0,0)$. In Section 4, we establish the existence of a heteroclinic connection between the equilibrium points $(\pm(1-p)^{1/(1-p)}, 0)$.

2 Self-Similar Structure

With $0 < p < 1$ and $T > 0$, we refer to $u : \mathbb{R} \times [0,T] \to \mathbb{R}$ as a solution to [CP] when u satisfies (1)-(2) with regularity,

$$u \in L^{\infty}(\mathbb{R} \times [0,T]) \cap C(\mathbb{R} \times [0,T]) \cap C^{2,1}(\mathbb{R} \times (0,T]). \tag{3}$$

Observe that $u^{\pm} : \mathbb{R} \times [0,T] \to \mathbb{R}$ given by

$$u^{\pm}(x,t) = \pm((1-p)t)^{1/(1-p)} \quad \forall (x,t) \in \mathbb{R} \times [0,T]$$

are the maximal and minimal solutions to [CP] (see [15, Chapter 8]), and hence any solution $u : \mathbb{R} \times [0,T] \to \mathbb{R}$ to [CP] must satisfy,

$$u^{-}(x,t) \leq u(x,t) \leq u^{+}(x,t) \quad \forall (x,t) \in \mathbb{R} \times [0,T]. \tag{4}$$

To construct spatially inhomogeneous solutions to [CP], we consider, for any fixed $x_0 \in \mathbb{R}$, self-similar solutions $u : \mathbb{R} \times [0,T] \to \mathbb{R}$ of the form,

$$u(x,t) = \begin{cases} w\left(\frac{x-x_0}{t^{1/2}}\right) t^{1/(1-p)} & , (x,t) \in \mathbb{R} \times (0,T], \\ 0 & , (x,t) \in \mathbb{R} \times \{0\}, \end{cases} \tag{5}$$

with $w : \mathbb{R} \to \mathbb{R}$ to be determined. Now, $u : \mathbb{R} \times [0, T] \to \mathbb{R}$ given by (5) is a solution to [CP] if and only if there exist constants α, $\beta \in \mathbb{R}$ such that $w : \mathbb{R} \to \mathbb{R}$ satisfies the following zero-value problem, namely,

$$w'' + \frac{1}{2}\eta w' + w|w|^{p-1} - \frac{1}{(1-p)}w = 0 \quad \forall \eta \in \mathbb{R}, \tag{6}$$

$$w(0) = \alpha, \quad w'(0) = \beta, \tag{7}$$

$$w \in C^2(\mathbb{R}) \cap L^\infty(\mathbb{R}). \tag{8}$$

Here $\eta = (x - x_0)/t^{1/2}$, and we observe that the ordinary differential equation (6) is both non-autonomous and non-Lipschitz. It is convenient to introduce

$$x = w, \quad y = w',$$

after which the problem (6)-(8) is equivalent to the zero-value problem for the two-dimensional, non-Lipschitz, non-autonomous, dynamical system,

$$x' = y \tag{9}$$

$$y' = \frac{1}{(1-p)}x - x|x|^{p-1} - \frac{1}{2}\eta y \quad \forall \eta \in \mathbb{R}, \tag{10}$$

$$(x(0), y(0)) = (\alpha, \beta), \tag{11}$$

$$(x, y) \in C^1(\mathbb{R}) \cap L^\infty(\mathbb{R}). \tag{12}$$

We refer to the equivalent zero-value problems in (6)-(8) and (9)-(12) as (S). Our objective is now to investigate those $(\alpha, \beta) \in \mathbb{R}^2$ for which (S) has a non-trivial solution. It is instructive to note, at this stage, via (4), that we may conclude that any solution to (S) must satisfy the inequality,

$$-(1-p)^{\frac{1}{(1-p)}} \leq w(\eta) \leq (1-p)^{\frac{1}{(1-p)}} \quad \forall \eta \in \mathbb{R}, \tag{13}$$

whilst, following [1, Corollary 2.6], any non-constant solution to (S) must be two-signed in w.

3 Homoclinic Connections

In this section we establish the existence of a two parameter family of homoclinic connections for (S) on the equilibrium point $(0, 0)$ of the dynamical system (9)-(10), and establish decay rates to the equilibrium point $(0, 0)$ as $|\eta| \to \infty$ on these homoclinic connections.

3.1 Existence

In this subsection, we establish the existence of homoclinic connections attached to the equilibrium point $(x, y) = (0, 0)$ of the dynamical system (9)-(10). To

begin, observe that $\mathbf{Q}:\mathbb{R}^3\to\mathbb{R}^2$, where

$$\mathbf{Q}(x,y,\eta)=(Q_1,Q_2)(x,y,\eta)=\left(y,\frac{1}{(1-p)}x-x|x|^{p-1}-\frac{1}{2}\eta y\right)\quad\forall(x,y,\eta)\in\mathbb{R}^3 \tag{14}$$

is such that $\mathbf{Q}\in C(\mathbb{R}^3)$, but also that $\mathbf{Q}$ is not locally Lipschitz continuous on $\mathbb{R}^3$ (note that $\mathbf{Q}$ is locally Lipschitz continuous on $\mathbb{R}^3\backslash N$, with N any neighbourhood of the plane $x=0$). We now have,

Theorem 1. *The problem (S) with zero-value $(\alpha,\beta)\in\mathbb{R}^2$ has a solution for $\eta\in[-\delta,\delta]$ (not necessarily unique), where $\delta=1/(1+M)$ and*

$$M=\max_{(x,y,\eta)\in R}|\mathbf{Q}(x,y,\eta)|$$

with

$$R=\{(x,y,\eta)\in\mathbb{R}^3:|x-\alpha|\le 1,\ |y-\beta|\le 1,\ |\eta|\le 1\}.$$

Proof. This follows immediately from the Cauchy-Peano Local Existence Theorem (see [6, Chapter 1, Theorem 1.2]) since $\mathbf{Q}:\mathbb{R}^3\to\mathbb{R}^2$ is such that $\mathbf{Q}\in C(\mathbb{R}^3)$. □

Remark 1. *When $\alpha\neq 0$, then the solution to (S) with zero-value $(\alpha,\beta)\in\mathbb{R}^2$ is unique for $\eta\in[-\delta',\delta']$ for some $0<\delta'\le\delta$. In addition, the problem (S) with zero-value $(\pm(1-p)^{1/(1-p)},0)$ has the unique global solution*

$$(x(\eta),y(\eta))=(\pm(1-p)^{1/(1-p)},0)\quad\forall\eta\in\mathbb{R}. \tag{15}$$

This follows since $\mathbf{Q}$ is locally Lipschitz in a neighbourhood of $(\pm(1-p)^{1/(1-p)},0)$ respectively. Also, the problem (S) with zero-value (0,0) has the unique global solution,

$$(x(\eta),y(\eta))=(0,0)\quad\forall\eta\in\mathbb{R}.$$

In this case uniqueness does not follows immediately, since $\mathbf{Q}$ is not locally Lipschitz continuous in any neighborhood of $(0,0)$, but instead follows after further qualitative results have been established for solutions to (S) (see Remark 2).

We now introduce the function $V:\mathbb{R}^2\to\mathbb{R}$ defined by,

$$V(x,y)=\frac{1}{2}y^2-\frac{1}{2(1-p)}x^2+\frac{1}{(1+p)}|x|^{1+p}\quad\forall(x,y)\in\mathbb{R}^2. \tag{16}$$

We observe immediately that

$$V\in C^{1,1}(\mathbb{R}^2), \tag{17}$$

with

$$\nabla V(x,y)=\left(\frac{-1}{(1-p)}x+x|x|^{p-1},y\right)\quad\forall(x,y)\in\mathbb{R}^2. \tag{18}$$



Figure 1: A qualitative sketch of the level curves of V

We now examine the structure of the level curves of V in $\mathbb{R}^2$, namely, the family of curves in $\mathbb{R}^2$ defined by

$$V(x,y) = c, \tag{19}$$

for $-\infty < c < \infty$. It is straightforward to establish that the family of level curves of V are qualitatively as sketched in Figure 1, with $\mathcal{H}$ representing the two level curves connecting $(-(1-p)^{1/1-p}, 0)$ to $((1-p)^{1/1-p}, 0)$ and enclosing the origin. In Figure 3.1, on the red curve $V = (1-p)^{2/(1-p)}/(2(1+p))$, whilst on the blue curves $V = 0$. At $(\pm(1-p)^{1/(1-p)}, 0)$ then $V = (1-p)^{2/(1-p)}/(2(1+p))$, whilst at $(0,0)$ then $V = 0$. Inside $\mathcal{H}$, the level curves are simple closed curves concentric with the origin $(0,0)$, and V is increasing from $V = 0$ at the origin $(0,0)$, as each level curve is crossed, when moving out from the origin $(0,0)$ to the boundary curve $\mathcal{H}$, on which $V = (1-p)^{2/(1-p)}/(2(1+p))$. Thus, inside $\mathcal{H}$, V has a minimum at the origin $(0,0)$ and is increasing on moving radially away from the origin $(0,0)$ to the boundary $\mathcal{H}$. On the level curves exterior and above or below $\mathcal{H}$, then $V > (1-p)^{2/(1-p)}/(2(1+p))$, whilst on the level curves to the left and right side of $\mathcal{H}$, then $V < (1-p)^{2/(1-p)}/(2(1+p))$, with $V = 0$ on the blue level curves. We now focus on the level curves of V on and inside $\mathcal{H}$, which have

$$0 \le c \le c^*(p), \tag{20}$$

where

$$c^*(p) = \frac{(1-p)^{2/(1-p)}}{2(1+p)}. \tag{21}$$

These are concentric closed curves surrounding the origin $(0,0)$. We will label the interior of the level curve $V = c$ by D_c, with the level curve $V = c$ labelled as ∂D_c, for $0 \le c \le c^*(p)$. In addition, we label the set

$$\bar{D}'_{c^*(p)} = \bar{D}_{c^*(p)} \backslash \{(\pm(1-p)^{1/(1-p)}, 0), (0,0)\}.$$

Now let $(x^*(\eta), y^*(\eta))$ be any solution to (S) for $\eta \in [-E, E]$ (any $E > 0$) with zero-value $(\alpha, \beta) \in \mathbb{R}^2$, and define $F : [-E, E] \to \mathbb{R}$ as,

$$F(\eta) = V(x^*(\eta), y^*(\eta)) \quad \forall \eta \in [-E, E]. \tag{22}$$

Then $F \in C^1([-E,E])$, and via (9), (10) and (14),

$$\begin{aligned} F'(\eta) =& \nabla V(x^*(\eta), y^*(\eta)).(x^{*'}(\eta), y^{*'}(\eta)) \\ =& \nabla V(x^*(\eta), y^*(\eta)).\mathbf{Q}(x^*(\eta), y^*(\eta), \eta) \quad \forall \eta \in [-E,E]. \end{aligned}$$

It then follows, via (18) and (14) that,

$$F'(\eta) = -\frac{1}{2}\eta(y^*(\eta))^2 \quad \forall \eta \in [-E,E]. \tag{23}$$

It follows from (23) that

$$F(\eta) \text{ is non-increasing for } \eta \in [0,E], \tag{24}$$

$$F(\eta) \text{ is non-decreasing for } \eta \in [-E,0]. \tag{25}$$

We can now establish the following,

Lemma 2. *Let $(x^*(\eta), y^*(\eta))$ be any solution to (S) on $[-E,E]$ (any $E > 0$) with zero-value $(\alpha,\beta) \in \bar{D}'_{c^*(p)}$. Then*

$$(x^*(\eta), y^*(\eta)) \in D_c \quad \forall \eta \in [-E,E]\backslash\{0\},$$

where $c = V(\alpha,\beta)$.

Proof. Let the zero-value $(\alpha,\beta) \in \partial D_c \backslash \{\pm((1-p)^{\frac{1}{1-p}}, 0)\}$ with

$$0 < c = V(\alpha,\beta) \leq c^*(p).$$

We first consider the case when $\beta \neq 0$. It follows from (23)-(25) that,

$$F(\eta) < F(0) \quad \forall \eta \in [-E,E]\backslash\{0\}. \tag{26}$$

Therefore, via (26),

$$V(x^*(\eta), y^*(\eta)) < c \quad \forall \eta \in [-E,E]\backslash\{0\},$$

and so

$$(x^*(\eta), y^*(\eta)) \in D_c \quad \forall \eta \in [-E,E]\backslash\{0\},$$

as required. Now consider the case when $\beta = 0$. Then $0 < |\alpha| < (1-p)^{1/(1-p)}$ and therefore, via (10) $y^{*'}(0) \neq 0$ after which a similar argument completes the proof. □

We now have:

Theorem 3. *For each $(\alpha,\beta) \in \bar{D}'_{c^*(p)}$, then (S) with zero-value (α,β) has a solution $(x^*(\eta), y^*(\eta))$ on $[-E,E]$ (any $E > 0$). Moreover, every such solution satisfies $(x^*(\eta), y^*(\eta)) \in D_c$ for all $\eta \in [-E,E]\backslash\{0\}$, where $c = V(\alpha,\beta)$.*

Proof. For any $(\alpha,\beta) \in \bar{D}'_{c^*(p)}$, Lemma 2 establishes that (S) with zero-value (α,β) is a priori bounded. The result then follows by a finite number of applications of the Cauchy-Peano Local Existence Theorem (see [6, Chapter 1, Theorem 1.2]), with $\delta = 1/(1+M)$ and

$$M = \max_{(x,y,\eta)\in R'} |\mathbf{Q}(x,y,\eta)|$$

whilst

$$R' = \{(x,y,\eta) \in \mathbb{R}^3 : |x| \le 2(1-p)^{1/(1-p)},\ |y| \le 2\sqrt{2c^*(p)},\ |\eta| \le 2E\}.$$

The final statement follows immediately from Lemma 2. □

We can now establish a global existence result for (S), namely

Corollary 4. *For $(\alpha,\beta) \in \bar{D}'_{c^*(p)}$ then (S) with zero-value (α,β) has a solution $(x^*(\eta),y^*(\eta))$ on $\mathbb{R}$. Moreover, every such solution satisfies $(x^*(\eta),y^*(\eta)) \in D_c$ for all $\eta \in \mathbb{R}\backslash\{0\}$, where $c = V(\alpha,\beta)$.*

Proof. Since Theorem 3 holds for any $E > 0$, the result follows immediately. □

Remark 2. *Let $(x^*(\eta),y^*(\eta))$ be any solution to (S) on $[-E,E]$ with zero-value $(0,0)$. It follows from (16), (22) and (23) that*

$$V(x^*(\eta),y^*(\eta)) = F(\eta) \le F(0) = V(0,0) = 0 \quad \forall \eta \in [-E,E]. \tag{27}$$

Thus $(x^*(\eta),y^*(\eta)) \in \mathcal{S}$ for all $\eta \in [-E,E]$, with $\mathcal{S}$ being a connected subset of

$$\{(x,y) \in \mathbb{R}^2 : V(x,y) \le 0\}$$

for which $(0,0) \in \mathcal{S}$. It follows that $\mathcal{S} = \{(0,0)\}$ and so $(x^*(\eta),y^*(\eta)) = (0,0)$ for all $\eta \in [-E,E]$. We conclude that the unique solution to (S) with zero-value $(0,0)$ is given by,

$$(x^*(\eta),y^*(\eta)) = (0,0) \quad \forall \eta \in \mathbb{R}.$$

We next introduce the function $H : \mathbb{R} \to \mathbb{R}$ such that

$$H(x) = \frac{1}{(1-p)}x - x|x|^{p-1} \quad \forall x \in \mathbb{R}, \tag{28}$$

and observe that

$$H \in C(\mathbb{R}). \tag{29}$$

We have,

Lemma 5. *Let $(\alpha,\beta) \in \bar{D}'_{c^*(p)}$, and let $(x^*(\eta),y^*(\eta))$ for $\eta \in \mathbb{R}$ be a global solution to (S) with zero-value (α,β). Then*

$$y^*(\eta) \to 0 \text{ as } |\eta| \to \infty.$$

Proof. We establish the result for $\eta \to \infty$; the result for $\eta \to -\infty$ follows similarly. Now, from (10),

$$y^{*'}(\eta) = H(x^*(\eta)) - \frac{1}{2}\eta y^*(\eta) \quad \forall \eta \in [0, \infty). \tag{30}$$

It then follows from (30) that,

$$y^*(\eta) = \beta e^{-\frac{1}{4}\eta^2} + e^{-\frac{1}{4}\eta^2} \int_0^\eta H(x^*(s)) e^{\frac{1}{4}s^2} ds \quad \forall \eta \in [0, \infty). \tag{31}$$

Thus,

$$|y^*(\eta)| \leq |\beta| e^{-\frac{1}{4}\eta^2} + e^{-\frac{1}{4}\eta^2} \int_0^\eta |H(x^*(s))| e^{\frac{1}{4}s^2} ds \quad \forall \eta \in [0, \infty). \tag{32}$$

However, via Corollary 4, $(x^*(\eta), y^*(\eta)) \in \bar{D}_{c^*(p)}$ for $\eta \in [0, \infty)$, and so, via (29), there exists a constant $M_H \geq 0$ such that

$$|H(x^*(s))| \leq M_H \quad \forall s \in [0, \infty). \tag{33}$$

It then follows from (32) and (33) that

$$|y^*(\eta)| \leq |\beta| e^{-\frac{1}{4}\eta^2} + M_H e^{-\frac{1}{4}\eta^2} \int_0^\eta e^{\frac{1}{4}s^2} ds \quad \forall \eta \in [0, \infty). \tag{34}$$

Now a simple application of Watson's Lemma (see [17, Proposition 2.1]), gives,

$$\int_0^\eta e^{\frac{1}{4}s^2} ds \sim \frac{2}{\eta} e^{\frac{1}{4}\eta^2} \text{ as } \eta \to \infty. \tag{35}$$

We then have, via (34) and (35), that

$$|y^*(\eta)| \leq |\beta| e^{-\frac{1}{4}\eta^2} + \frac{4M_H}{\eta} \text{ as } \eta \to \infty. \tag{36}$$

It follows from (36) that

$$y^*(\eta) \to 0 \text{ as } \eta \to \infty,$$

as required □

We next have,

Lemma 6. *Let $(x^*(\eta), y^*(\eta))$ for $\eta \in \mathbb{R}$ be a global solution to (S) with zero-value $(\alpha, \beta) \in \bar{D}'_{c^*(p)}$, and $F : \mathbb{R} \to \mathbb{R}$ as in (22). Then $F(\eta)$ is non-increasing for $\eta \in (0, \infty)$ and non-decreasing for $\eta \in (-\infty, 0)$, with*

$$F(\eta) \to \begin{cases} F_\infty & \text{as } \eta \to \infty \\ F_{-\infty} & \text{as } \eta \to -\infty \end{cases}$$

where $F_\infty, F_{-\infty} \in [0, F(0))$.

Proof. We observe from Corollary 4 that

$$(x^*(\eta), y^*(\eta)) \in D_c \quad \forall \eta \in \mathbb{R}\backslash\{0\}, \tag{37}$$

with $c = V(\alpha, \beta) = F(0)$, and so,

$$0 \leq F(\eta) < F(0) \quad \forall \eta \in \mathbb{R}\backslash\{0\}. \tag{38}$$

In addition, it follows from (38), (24) and (25), since $F \in C^1(\mathbb{R})$, that there exist $F_\infty, F_{-\infty} \in \mathbb{R}$, such that

$$F(\eta) \to \begin{cases} F_\infty & \text{as } \eta \to \infty \\ F_{-\infty} & \text{as } \eta \to -\infty \end{cases}$$

where $F_\infty, F_{-\infty} \in [0, F(0))$, as required. □

We now have,

Theorem 7. *Let $(x^*(\eta), y^*(\eta))$ for $\eta \in \mathbb{R}$ be a global solution to (S) with zero-value $(\alpha, \beta) \in \bar{D}'_{c^*(p)}$. Then,*

$$(x^*(\eta), y^*(\eta)) \to (0, 0) \text{ as } |\eta| \to \infty.$$

Proof. We establish the result for $\eta \to \infty$. The result for $\eta \to -\infty$ follows similarly. We first recall from Corollary 4 that,

$$(x^*(\eta), y^*(\eta)) \in D_{c^*(p)} \quad \forall \eta \in \mathbb{R}\backslash\{0\}, \tag{39}$$

and from Lemma 5 that,

$$y^*(\eta) \to 0 \text{ as } \eta \to \infty. \tag{40}$$

In addition, we have from Lemma 6 that,

$$V(x^*(\eta), y^*(\eta)) \to F_\infty \text{ as } \eta \to \infty \tag{41}$$

for some $F_\infty \in [0, c^*(p))$. It follows from (39)-(41) that

$$x^*(\eta) \to x_\infty \text{ or } x^*(\eta) \to -x_\infty \text{ as } \eta \to \infty \tag{42}$$

where x_∞ is the single non-negative root of

$$V(x, 0) = F_\infty \text{ with } x \in [0, (1-p)^{1/(1-p)}).$$

Without loss of generality we will suppose that

$$(x^*(\eta), y^*(\eta)) \to (x_\infty, 0) \text{ as } \eta \to \infty. \tag{43}$$

However, it follows from (10) that,

$$y^*(\eta) = \beta e^{-\frac{1}{4}\eta^2} + e^{-\frac{1}{4}\eta^2} \int_0^\eta H(x^*(s)) e^{\frac{1}{4}s^2} ds \quad \eta \in [0, \infty) \tag{44}$$

with $H : \mathbb{R} \to \mathbb{R}$ given by (28), and

$$H(x_\infty) \leq 0. \tag{45}$$

Using (42), it is straightforward to establish that, when,

$$H(x_\infty) < 0, \tag{46}$$

then from (44),

$$y^*(\eta) \sim \frac{2H(x_\infty)}{\eta} \text{ as } \eta \to \infty. \tag{47}$$

In addition, from (9), we have,

$$x^*(\eta) = \alpha + \int_0^\eta y^*(s)ds \quad \forall \eta \in [0, \infty), \tag{48}$$

which gives, via (47), that

$$x^*(\eta) \sim 2H(x_\infty) \log \eta, \quad \text{as } \eta \to \infty,$$

which contradicts (42). We conclude that (46) cannot hold, and so, via (45), we must have

$$H(x_\infty) = 0, \tag{49}$$

which, since $x_\infty \in [0, (1-p)^{1/(1-p)})$, requires $x_\infty = 0$. It then follows from (43) that,

$$(x^*(\eta), y^*(\eta)) \to (0, 0) \text{ as } \eta \to \infty,$$

as required. □

We conclude from Corollary 4 and Theorem 7 that the problem (S) has a two parameter family of nontrivial, distinct homoclinic connections on the equilibrium point $(0, 0)$, parametrized by $(\alpha, \beta) \in \bar{D}'_{c^*(p)}$ which we will denote by $w_{\alpha,\beta} : \mathbb{R} \to \mathbb{R}$ for each $(\alpha, \beta) \in \bar{D}'_{c^*(p)}$. Here $w = w_{\alpha,\beta}(\eta)$, $\eta \in \mathbb{R}$, has zero-values $w(0) = \alpha$, $w'(0) = \beta$. Moreover,

$$(w_{\alpha,\beta}(\eta), w'_{\alpha,\beta}(\eta)) \in D_{V(\alpha,\beta)} \quad \forall \eta \in \mathbb{R} \backslash \{0\}.$$

Additionally, note that $w_{0,\beta}(\eta)$ is an odd function of η whilst $w_{\alpha,0}(\eta)$ is an even function of η. Furthermore, it also follows from the comments below (13) that $w_{\alpha,\beta}(\eta)$ must be two signed for $\eta \in \mathbb{R}$.

3.2 Decay Bounds and Estimates

In this section, we establish results concerning the rate of decay to zero of $w_{\alpha,\beta}(\eta)$ as $\eta \to \pm\infty$. Specifically, we establish algebraic bounds on the rate of decay of $w_{\alpha,\beta}(\eta)$ as $\eta \to \pm\infty$, and hence, determine that $w_{\alpha,\beta} \in L_q(\mathbb{R})$ for each $q > (1-p)/2$. From these bounds we may infer that the corresponding solution to [CP], say $u_{\alpha,\beta} : \mathbb{R} \times [0, \infty) \to \mathbb{R}$, satisfies $u(\cdot, t) \in L_q(\mathbb{R})$ for each $t \in [0, \infty)$ and

$q > (1-p)/2$. To complement the algebraic bounds, we also provide a rational asymptotic approximation to the decay rate of $w_{\alpha,\beta}(\eta)$ as $\eta \to \pm\infty$, which, in fact suggests exponential decay as $\eta \to \pm\infty$.

To begin, observe that $w = w_{\alpha,\beta}(\eta)$ for $\eta \in \mathbb{R}$, via (6), satisfies

$$(e^{\frac{1}{4}\eta^2} w')' = H(w) e^{\frac{1}{4}\eta^2} \quad \forall \eta \in \mathbb{R}.$$

It follows from two successive integrations, that

$$w'(\eta) = \beta e^{-\frac{1}{4}\eta^2} + e^{-\frac{1}{4}\eta^2} \int_0^\eta H(w(s)) e^{\frac{1}{4}s^2} ds \quad \forall \eta \in \mathbb{R} \tag{50}$$

whilst,

$$w(\eta) = \alpha + \int_0^\eta \beta e^{-\frac{1}{4}t^2} dt + \int_0^\eta e^{-\frac{1}{4}t^2} \int_0^t H(w(s)) e^{\frac{1}{4}s^2} ds dt \quad \forall \eta \in \mathbb{R}. \tag{51}$$

We now have,

Proposition 8. *Let $w : \mathbb{R} \to \mathbb{R}$ be a solution to (S) with zero-value $(\alpha, \beta) \in \bar{D}'_{c^*(p)}$. Suppose that*

$$|w(\eta)| \leq \frac{c_1}{(1+|\eta|)^\sigma} \quad \forall \eta \in \mathbb{R}.$$

with $\sigma \geq 0$ and $c_1 > 0$ (independent of α and β). Then, there exists $c_2 > 0$, which depends on c_1, σ and p, (independent of α and β) such that,

$$|w'(\eta)| \leq \frac{c_2}{(1+|\eta|)^{\sigma p+1}} \quad \forall \eta \in \mathbb{R}.$$

Proof. We give a proof for $\eta \geq 0$; the result for $\eta < 0$ follows similarly. Observe that

$$|H(w(\eta))| = \left| \frac{1}{(1-p)} w(\eta) - |w(\eta)|^{p-1} w(\eta) \right| \leq \frac{c_1^p}{(1+\eta)^{\sigma p}} \quad \forall \eta \in [0, \infty), \tag{52}$$

since, via Corollary 4, $|w(\eta)| < (1-p)^{\frac{1}{(1-p)}}$ for $\eta \in [0, \infty)$. Thus, via (50) and (52), we have,

$$|w'(\eta)| \leq |\beta| e^{-\frac{1}{4}\eta^2} + c_1^p e^{-\frac{1}{4}\eta^2} \int_0^\eta \frac{1}{(1+s)^{\sigma p}} e^{\frac{1}{4}s^2} ds \quad \forall \eta \in [0, \infty). \tag{53}$$

Now, the second term on the right hand side of (53) is a non-negative continuous function for $\eta \in [0, \infty)$, with asymptotic form,

$$c_1^p e^{-\frac{1}{4}\eta^2} \int_0^\eta \frac{1}{(1+s)^{\sigma p}} e^{\frac{1}{4}s^2} ds \sim \frac{2c_1^p}{\eta^{\sigma p+1}} \text{ as } \eta \to \infty.$$

It follows that,

$$c_1^p e^{-\frac{1}{4}\eta^2} \int_0^\eta \frac{1}{(1+s)^{\sigma p}} e^{\frac{1}{4}s^2} ds \leq \frac{4c_1^p}{\eta^{\sigma p+1}} \text{ as } \eta \to \infty.$$

We conclude that there exists a positive constant c_2, depending upon c_1, p, and σ, such that

$$|w'(\eta)| \leq \frac{c_2}{(1+\eta)^{\sigma p+1}} \quad \forall \eta \in [0,\infty),$$

as required. □

We next have,

Proposition 9. *Let $w : \mathbb{R} \to \mathbb{R}$ be a solution to (S) with zero-value $(\alpha,\beta) \in \bar{D}'_{c^*(p)}$. Then,*

$$(w(\eta), w'(\eta)) \to (0,0) \text{ as } \eta \to \pm\infty,$$

and moreover,

$$|w'(\eta)| \leq \frac{c_2}{(1+|\eta|)} \quad \forall \eta \in \mathbb{R},$$

with $c_2 > 0$ dependent upon p (independent of α and β).

Proof. The first conclusion follows directly from Theorem 7. Additionally, it follows from Corollary 4 that

$$(w(\eta), w'(\eta)) \in \mathcal{H} \quad \forall \eta \in \mathbb{R},$$

and hence, it follows from Proposition 8 (with $\sigma = 0$, $c_1 = (1-p)^{1/(1-p)}$) that

$$|w'(\eta)| \leq \frac{c_2}{(1+|\eta|)} \quad \forall \eta \in \mathbb{R},$$

as required. □

We now demonstrate that every solution $w : \mathbb{R} \to \mathbb{R}$ to (S) with zero-value $(\alpha,\beta) \in \bar{D}'_{c^*(p)}$ decays to zero as $\eta \to \pm\infty$, with decay rate which is at least algebraic in η as $\eta \to \pm\infty$. In particular, we demonstrate that $w : \mathbb{R} \to \mathbb{R}$ is contained in $L^q(\mathbb{R})$ for any $q > (1-p)/2$. The proof is based on the decay bounds obtained in [10].

Theorem 10. *Let $w : \mathbb{R} \to \mathbb{R}$ be a solution to (S) with zero-value $(\alpha,\beta) \in \bar{D}'_{c^*(p)}$. Then, for any $\epsilon > 0$, there exists $c_{1\epsilon}, c_{2\epsilon} > 0$ (dependent generally on α, β, p and ϵ) such that*

$$|w(\eta)| < \frac{c_{1\epsilon}}{(1+|\eta|)^{\frac{2}{(1-p)}-\epsilon}} \quad \forall \eta \in \mathbb{R},$$

$$|w'(\eta)| < \frac{c_{2\epsilon}}{(1+|\eta|)^{\frac{(1+p)}{(1-p)}-\epsilon}} \quad \forall \eta \in \mathbb{R}.$$

Proof. We give a proof for $\eta \geq 0$; the argument for $\eta < 0$ follows similarly. Observe on multiplying (6) by $\eta^{-1}w(\eta)$, we have,

$$\frac{1}{\eta}\left[|w(\eta)|^{1+p} - \frac{(w(\eta))^2}{(1-p)}\right] = -\left[\frac{(w(\eta))^2}{4} + \frac{w(\eta)w'(\eta)}{\eta}\right]' + \frac{(w'(\eta))^2}{\eta} - \frac{w(\eta)w'(\eta)}{\eta^2} \tag{54}$$

for $\eta \in (0,\infty)$. Additionally, via Proposition 9, it follows that there exists $\eta_* \in (0,\infty)$ such that,

$$|w(\eta)| \leq \left(\frac{2p(1-p)}{(1+p)}\right)^{\frac{1}{(1-p)}} \qquad \forall \eta \in [\eta_*, \infty), \tag{55}$$

and for $F : [0,\infty) \to \mathbb{R}$ given by

$$F(\eta) = V(w(\eta), w'(\eta)) \quad \forall \eta \in [0,\infty),$$

that

$$0 \leq F(\eta) \leq \left(\frac{4(c(p))^{\frac{2}{(1+p)}}}{C(p)}\right)^{(1+p)/(1-p)} \qquad \eta \in [\eta_*, \infty), \tag{56}$$

where

$$c(p) = \frac{1}{(1+p)} - \frac{1}{2}, \text{ and } C(p) = \frac{2(1+p)}{(1-p)} + 1. \tag{57}$$

Thus, it follows from (54) that

$$\begin{aligned}
\frac{F(\eta)}{\eta} &= \frac{(w'(\eta))^2}{2\eta} + \frac{1}{\eta}\left[-\frac{(w(\eta))^2}{2(1-p)} + \frac{|w(\eta)|^{1+p}}{(1+p)}\right] \\
&\leq \frac{(w'(\eta))^2}{2\eta} + \frac{1}{\eta}\left[-\frac{(w(\eta))^2}{(1-p)} + |w(\eta)|^{1+p}\right] \\
&= \frac{3(w'(\eta))^2}{2\eta} - \left[\frac{(w(\eta))^2}{4} + \frac{w(\eta)w'(\eta)}{\eta}\right]' - \frac{w(\eta)w'(\eta)}{\eta^2},
\end{aligned} \tag{58}$$

for $\eta \in [\eta_*, \infty)$. Since $F(\eta) \geq 0$ for all $\eta \in [\eta^*, \infty)$, together with the decay estimates in Proposition 9, it follows that we may integrate inequality (58) from η ($\geq \eta^*$) to l, and then allow $l \to \infty$, to obtain,

$$\int_\eta^\infty \frac{F(t)}{t} dt \leq \frac{(w(\eta))^2}{4} + \frac{2}{\eta} \sup_{t \geq \eta} |w(t)w'(t)| + \frac{3}{2}\int_\eta^\infty \frac{(w'(t))^2}{t} dt \tag{59}$$

for $\eta \in [\eta_*, \infty)$. We also note, that since, via Corollary 4, $|w(\eta)| < (1-p)^{1/(1-p)}$, we have,

$$F(\eta) \geq |w(\eta)|^{1+p} c(p) \geq 0, \tag{60}$$

for $\eta \in [\eta_*, \infty)$. It therefore follows from (59) and (60) that

$$0 \leq \int_\eta^\infty \frac{F(t)}{t} dt \leq \frac{1}{4}\left(\frac{F(\eta)}{c(p)}\right)^{\frac{2}{(1+p)}} + \frac{2}{\eta} \sup_{t \geq \eta} |w(t)w'(t)| + \frac{3}{2}\int_\eta^\infty \frac{(w'(t))^2}{t} dt \tag{61}$$

for $\eta \in [\eta_*, \infty)$. We observe that the right hand side of (61) is uniformly bounded for $\eta \in [\eta_*, \infty)$ via Proposition 9.

Now suppose that there exists $k > 0$ such that

$$F(\eta) \leq \frac{k}{\eta^\sigma} \quad \forall \eta \in [\eta_*, \infty) \tag{62}$$

for some $\sigma \geq 0$ (note that (62) holds when $\sigma = 0$ via Proposition 9). Then, via (60), it follows that there exists $c_1 > 0$ such that

$$|w(\eta)| \leq \frac{c_1}{\eta^{\frac{\sigma}{(1+p)}}} \quad \forall \eta \in [\eta_*, \infty) \tag{63}$$

and so, via Proposition 8, there exists $c_2 > 0$ such that

$$|w'(\eta)| \leq \frac{c_2}{\eta^{\frac{\sigma p}{(1+p)}+1}} \quad \forall \eta \in [\eta_*, \infty). \tag{64}$$

Thus, it follows from (61)-(64) and (56), that there exists $c_3, c_4, c_5 > 0$ such that

$$\begin{aligned} \int_\eta^\infty \frac{F(t)}{t} dt &\leq \frac{1}{4}\left(\frac{F(\eta)}{c(p)}\right)^{\frac{2}{(1+p)}} + \frac{c_3}{\eta^{\sigma+2}} + \frac{c_4}{\eta^{\frac{2\sigma p}{(1+p)}+2}} \\ &\leq \frac{F(\eta)}{C(p)} + \frac{c_5}{\eta^{\frac{2\sigma p}{(1+p)}+2}} \end{aligned} \tag{65}$$

for $\eta \in [\eta_*, \infty)$. Upon setting $G : [\eta_*, \infty) \to \mathbb{R}$ to be

$$G(\eta) = \int_\eta^\infty \frac{F(t)}{t} dt \quad \forall \eta \in [\eta_*, \infty),$$

it follows from (65) that G satisfies,

$$\left(t^{C(p)} G(t)\right)' \leq \frac{c_6}{t^{\frac{2\sigma p}{(1+p)}+3-C(p)}} \quad \forall t \in [\eta_*, \infty), \tag{66}$$

with $c_6 > 0$ constant. An integration of (66) gives

$$G(\eta) \leq \frac{c_7}{\eta^{\frac{2\sigma p}{(1+p)}+2}} + \frac{c_8}{\eta^{C(p)}} \quad \forall \eta \in [\eta_*, \infty), \tag{67}$$

with $c_7, c_8 > 0$ constants. Also, recalling, via Lemma 6, that $F(\eta)$ is non-increasing on $[\eta^*, \infty)$, we have,

$$G(\eta) \geq \int_\eta^{2\eta} \frac{F(t)}{t} dt \geq \frac{1}{2} F(2\eta), \quad \forall \eta \in [\eta_*, \infty). \tag{68}$$

Thus, it follows from (67) and (68) that there exist constants $c_9, c_{10} > 0$ such that

$$F(\eta) \leq \frac{c_9}{\eta^{\frac{2\sigma p}{(1+p)}+2}} + \frac{c_{10}}{\eta^{C(p)}} \quad \forall \eta \in [\eta_*, \infty). \tag{69}$$

Since (62) holds for $\sigma = 0$, it follows that there exists sequences $\{\sigma_n\}_{n\in\mathbb{N}}$ and $\{k_n\}_{n\in\mathbb{N}}$ given by

$$\sigma_1 = 0, \quad \sigma_{n+1} = \min\left\{\frac{2\sigma_n p}{(1+p)} + 2,\ C(p)\right\} \tag{70}$$

such that

$$F(\eta) \leq \frac{k_n}{\eta^{\sigma_n}} \quad \forall \eta \in [\eta_*, \infty). \tag{71}$$

We obtain from (70) and (57) that,

$$\sigma_n = \frac{2(1+p)}{(1-p)} - \frac{4p}{(1-p)}\left(\frac{2p}{(1+p)}\right)^{n-2} \qquad \forall n \in \mathbb{N}$$

and hence σ_n is increasing with

$$\sigma_n \to \frac{2(1+p)}{(1-p)} \quad \text{as } n \to \infty. \tag{72}$$

Therefore it follows, via (60) and (70)-(72), that for each $\epsilon > 0$ there exists $c_{1\epsilon} > 0$ such that

$$|w(\eta)| \leq \frac{c_{1\epsilon}}{(1+\eta)^{\frac{2}{(1-p)}-\epsilon}} \quad \forall \eta \in [0, \infty), \tag{73}$$

recalling that $w(\eta)$ is bounded on $[0, \eta^*]$. The bound on $|w'(\eta)|$ follows immediately from (73) and Proposition 8. □

The algebraic bounds in Theorem 10 are the tightest decay rates we have been able to establish rigorously. However, the following asymptotic argument indicates that, in fact, $w = w_{\alpha,\beta}(\eta)$ decays exponentially in η as $|\eta| \to \infty$, accompanied by rapid oscillatory behaviour. To this end, we now consider the asymptotic structure of $w = w_{\alpha,\beta}(\eta)$ as $\eta \to \infty$, with the same structure following as $\eta \to -\infty$. Now, for $\eta >> 1$, then $w = w_{\alpha,\beta}(\eta)$ satisfies,

$$w'' + \frac{1}{2}\eta w' + w|w|^{p-1} - \frac{1}{(1-p)}w = 0 \quad \eta >> 1 \tag{74}$$

$$w(\eta),\ w'(\eta) \to 0 \quad \text{as } \eta \to \infty, \tag{75}$$

via (6) and Proposition 9. On using (75), the dominant form of (74) when $\eta >> 1$ is

$$w'' + w|w|^{p-1} = 0. \tag{76}$$

Every solution to (76) is periodic and may be written (up to translation in η) as,

$$w(\eta, a) = aW\left(a^{-\frac{1}{2}(1-p)}\eta\right), \quad \forall \eta \in \mathbb{R}, \tag{77}$$

where $a \in \mathbb{R}^+$ is a parameter and $W : \mathbb{R} \to \mathbb{R}$ is that unique periodic function which satisfies the problem,

$$W'' + W|W|^{p-1} = 0, \quad \zeta \in \mathbb{R} \tag{78}$$

$$W(0) = 1, \quad W'(0) = 0. \tag{79}$$

The period of $W(\zeta)$ is given by

$$T(p) = 2^{3/2}(1+p)^{1/2}\int_0^1 \frac{d\lambda}{(1-\lambda^{(1+p)})^{1/2}} \tag{80}$$



Figure 2: Phase paths for solutions to (78)-(79) for $p_k = (0.1)k$ for $k = 1...9$. Here the phase path for p_k encloses the phase path for p_{k+1} for $k = 1...8$.

whilst,

$$W(\zeta) = -W\left(\frac{1}{2}T(p) - \zeta\right) = W(-\zeta) \quad \forall \zeta \in \mathbb{R}. \tag{81}$$

Via an integration, the solution to (78)-(79) satisfies

$$\frac{(W'(\eta))^2}{2} + \frac{|W(\eta)|^{1+p}}{(1+p)} = \frac{1}{(1+p)} \quad \forall \eta \in \mathbb{R},$$

which represents a periodic orbit in the (W, W') phase plane, as illustrated in Figure 3.2. It follows from (77) that $w(\eta, a)$ has amplitude $a > 0$ and period

$$T_a(p) = a^{\frac{1}{2}(1-p)}T(p). \tag{82}$$

For any fixed $a \in \mathbb{R}^+$, (77) cannot represent the asymptotic structure to (74) and (75) since W is periodic. The remaining terms in (74) must induce decay as $\eta \to \infty$. However, we observe from (82) that the oscillations in $w(\eta, a)$ becomes increasingly rapid as the amplitude $a \to 0^+$. This suggests that we seek the asymptotic structure of (74)-(75) as $\eta \to \infty$ in the form,

$$w(\eta) \sim a(\eta)W(a(\eta)^{-\frac{1}{2}(1-p)}\eta) \text{ as } \eta \to \infty, \tag{83}$$

with $a(\eta) > 0$ and,

$$a(\eta), a'(\eta) \to 0 \text{ as } \eta \to \infty. \tag{84}$$

Now, the rate of change of amplitude of oscillation in (83), $a'(\eta)$, approaches zero as $\eta \to \infty$, whilst the frequency of oscillation becomes unbounded as $\eta \to \infty$. We can thus use an averaging approach to determine an evolution equation for the amplitude $a(\eta)$ as $\eta \to \infty$. We substitute (83) into (6) and make use of (78).

We then integrate the resulting ordinary differential equation over *one* period of $W(\cdot)$, over which, we may hold a fixed. We obtain the leading order amplitude equation as,

$$a'' + \frac{1}{2}\eta a' - \frac{1}{(1-p)}a = 0, \quad \eta >> 1, \tag{85}$$

$$a(\eta), a'(\eta) \to 0 \text{ as } \eta \to \infty. \tag{86}$$

The linear ordinary differential equation (85) has two basis functions $a_+ : \mathbb{R} \to \mathbb{R}$ and $a_- : \mathbb{R} \to \mathbb{R}$ which have

$$a_+(\eta) \sim \eta^{-\left(1+\frac{2}{(1-p)}\right)} e^{-\frac{1}{4}\eta^2}, \quad a_-(\eta) \sim \eta^{\frac{2}{(1-p)}} \quad \text{as } \eta \to \infty.$$

It follows that

$$a(\eta) \sim A_\infty \eta^{-\left(1+\frac{2}{(1-p)}\right)} e^{-\frac{1}{4}\eta^2} \quad \text{as } \eta \to \infty, \tag{87}$$

with A_∞ being a positive globally determined constant dependent, in general, on α, β and p. Thus, from (83), we have

$$w_{\alpha,\beta}(\eta) \sim a(\eta) W(a(\eta)^{-\frac{1}{2}(1-p)}\eta) \text{ as } \eta \to \infty, \tag{88}$$

with, $\alpha(\eta)$ having the asymptotic form (87) as $\eta \to \infty$. The same argument leads to the same (up to the constant A_∞) asymptotic structure as $\eta \to -\infty$. As a consequence of (87) and (88), we anticipate that $w_{\alpha,\beta}(\eta)$ decays to zero at a Gaussian rate as $|\eta| \to \infty$, whilst oscillating about zero with a local frequency which increases at a Gaussian rate as $|\eta| \to \infty$. This indicates that, in fact, $w_{\alpha,\beta} \in L^q(\mathbb{R})$ for any $q > 0$.

3.3 Localized Solutions to [CP]

Following Corollary 4 and Theorem 7, for each $(\alpha, \beta) \in \bar{D}'_{c^*(p)}$, we have constructed a non-trivial, localized, global solution $u_{\alpha,\beta} : \mathbb{R} \times [0, \infty) \to \mathbb{R}$ to [CP], namely,

$$u_{\alpha,\beta}(x,t) = \begin{cases} t^{\frac{1}{(1-p)}} w_{\alpha,\beta}\left(\frac{x}{t^{1/2}}\right) & , (x,t) \in \mathbb{R} \times (0,\infty) \\ 0 & , (x,t) \in \mathbb{R} \times \{0\}. \end{cases} \tag{89}$$

With this two parameter family of solutions to [CP], each solution is distinct, and is not a spatial translate of any other solution in the family. However, we observe that $u_{\alpha,\beta}(x - x_0, t)$ is also a global solution to [CP] for any fixed $x_0 \in \mathbb{R}$. A trivial calculation from (89) establishes that

$$(u_{\alpha,\beta})_x(x,t) = t^{\frac{1}{(1-p)}-\frac{1}{2}} w'_{\alpha,\beta}\left(\frac{x}{t^{1/2}}\right), \tag{90}$$

$$(u_{\alpha,\beta})_t(x,t) = \frac{1}{(1-p)} t^{\frac{1}{(1-p)}-1}\left(w_{\alpha,\beta}\left(\frac{x}{t^{1/2}}\right) - \frac{1}{2}(1-p)\left(\frac{x}{t^{1/2}}\right) w'_{\alpha,\beta}\left(\frac{x}{t^{1/2}}\right)\right), \tag{91}$$

for $(x,t) \in \mathbb{R} \times (0,\infty)$, whilst from (1),

$$(u_{\alpha,\beta})_{xx}(x,t) = (u_{\alpha,\beta})_t(x,t) - (u_{\alpha,\beta}|u_{\alpha,\beta}|^{p-1})(x,t), \tag{92}$$

for $(x,t) \in \mathbb{R} \times (0,\infty)$. It then follows immediately from Theorem 7 that,

$$(u_{\alpha,\beta})_x,\ (u_{\alpha,\beta})_t,\ (u_{\alpha,\beta})_{xx} \to 0 \text{ as } t \to 0^+ \text{ uniformly for } x \in \mathbb{R},$$

and so, in fact,

$$u_{\alpha,\beta} \in L^\infty(\mathbb{R} \times [0,T]) \cap C(\mathbb{R} \times [0,T]) \cap C^{2,1}(\mathbb{R} \times [0,T]). \tag{93}$$

It follows from (93) that for each $(\alpha,\beta) \in \bar{D}'_{c^*(p)}$, and any $\tau > 0$, then $u^\tau_{\alpha,\beta} : \mathbb{R} \times [0,\infty) \to \mathbb{R}$ such that

$$u^\tau_{\alpha,\beta}(x,t) = \begin{cases} (t-\tau)^{\frac{1}{(1-p)}} w_{\alpha,\beta}\left(\frac{x}{(t-\tau)^{1/2}}\right) & ,(x,t) \in \mathbb{R} \times (\tau,\infty) \\ 0 & ,(x,t) \in \mathbb{R} \times [0,\tau] \end{cases}$$

is also a non-trivial, localized, global solution to [CP]. Finally, we observe, via Theorem 10 that for each $q > (1-p)/2$, then $u_{\alpha,\beta}(\cdot,t) \in L^q(\mathbb{R}$ for each $t \geq 0$. Moreover, (87) and (88) suggest that the localization is Gaussian in x for each $t > 0$.

4 Heteroclinic Connections

In this section we establish the existence of at least one heteroclinic connection for (S) from the equilibrium point $(-(1-p)^{1/(1-p)}, 0)$ to the equilibrium point $((1-p)^{1/(1-p)}, 0)$.

4.1 Existence

We first consider solutions to the problem (S) for $\eta \in [0,\infty)$ and which remain in the region $\Omega \subset \mathbb{R}^2$, given as

$$\Omega = \left\{(x,y) : 0 < x < (1-p)^{1/(1-p)}, y > 0\right\} \tag{94}$$

with boundary $\partial\Omega = \overline{\Omega} \setminus \Omega$. We also define the following subset of $\partial\Omega$, namely,

$$\partial\Omega_1 = \{(x,y) : x = 0, y > 0\}. \tag{95}$$

Specifically, we consider (S) for $\eta \in [0,\infty)$ and demonstrate that there exists a solution $(x,y) : [0,\infty) \to \overline{\Omega}$ with zero-value $(0,\beta) \in \partial\Omega_1$ and which satisfies

$$(x(\eta), y(\eta)) \in \Omega \quad \forall \eta \in (0,\infty), \tag{96}$$

$$(x(\eta), y(\eta)) \to ((1-p)^{1/(1-p)}, 0) \quad \text{as } \eta \to \infty. \tag{97}$$

To begin with, it is readily established that for each zero-value $(0,\beta) \in \partial\Omega_1$, then (S) has a local solution $(x,y):[0,\delta]\to\mathbb{R}^2$ (for some $\delta>0$). Moreover, $(x(\eta),y(\eta))\in\Omega$ for $\eta\in(0,\delta]$, and $x(\eta)$ is monotone increasing whilst $y(\eta)$ is monotone decreasing, with $\eta\in(0,\delta]$. It is then straightforward to establish that $(x(\eta),y(\eta))$ can be *uniquely* continued beyond $\eta=\delta$ and must satisfy one of the following three possibilities:

(i) There exists $\eta_\beta>0$ such that $(x(\eta),y(\eta))\in\Omega$ for all $\eta\in(0,\eta_\beta)$ and $(x(\eta_\beta),y(\eta_\beta))=((1-p)^{\frac{1}{1-p}},y_\beta)$ with $0<y_\beta<\beta$, whilst $x'(\eta_\beta)=y_\beta>0$, and so there exists $\epsilon_\beta>0$ such that $(x(\eta),y(\eta))\notin\bar{\Omega}\cup(\{0\}\times\mathbb{R})$ for $\eta\in(\eta_\beta,\eta_\beta+\epsilon_\beta]$.

(ii) There exists $\eta_\beta>0$ such that $(x(\eta),y(\eta))\in\Omega$ for all $\eta\in(0,\eta_\beta)$ and $(x(\eta_\beta),y(\eta_\beta))=(x_\beta,0)$ with $0<x_\beta<(1-p)^{\frac{1}{1-p}}$, whilst $y'(\eta_\beta)<0$ and so there exists $\epsilon_\beta>0$ such that $(x(\eta),y(\eta))\notin\bar{\Omega}\cup(\{0\}\times\mathbb{R})$ for $\eta\in(\eta_\beta,\eta_\beta+\epsilon_\beta]$.

(iii) $(x(\eta),y(\eta))\in\Omega$ for all $\eta\in(0,\infty)$ and $(x(\eta),y(\eta))\to((1-p)^{\frac{1}{1-p}},0)$ as $\eta\to\infty$.

Our aim now is to obtain a uniqueness result for (S) with zero-value in $\partial\Omega_1$, and from this a continuous dependence result. This is non-trivial, since $\mathbf{Q}$ in (14) is not locally Lipschitz continuous in any neighborhood of $(0,\beta)\in\partial\Omega_1$, and so standard uniqueness and continuous dependence theory fail to apply. To begin with, we provide a local a priori bound for any solution of (S) with zero-value $(0,\beta)\in\partial\Omega_1$.

Proposition 11. *Let $(x,y):[0,\eta_\beta]\to\mathbb{R}^2$ be any solution to (S) with zero-value $(0,\beta)\in\partial\Omega_1$ and which satisfies either case (i) or (ii). Then,*

$$\eta_\beta>\min\left\{\left(\frac{2}{\beta}\right)\left(m_H+\sqrt{m_H^2+\frac{(\beta)^2}{2}}\right),\ \frac{(1-p)^{1/(1-p)}}{\beta}\right\}=\eta^* \tag{98}$$

with

$$m_H=\inf_{\lambda\in[0,(1-p)^{1/(1-p)}]}H(\lambda), \tag{99}$$

Proof. Let $(x,y):[0,\eta_\beta]\to\mathbb{R}^2$ be any solution to (S) with zero-value $(0,\beta)\in\partial\Omega_1$, and which satisfies either case (i) or case (ii). Suppose that $\eta_\beta\le\eta^*$. Since $(x(\eta),y(\eta))\in\Omega$ for all $\eta\in(0,\eta_\beta)$, it follows from (10) that

$$\beta+m_H\eta-\frac{\beta}{4}\eta^2<y(\eta)<\beta\quad\forall\eta\in(0,\eta_\beta]. \tag{100}$$

However, $\eta_\beta\le\eta^*$ and so, via (100),

$$\frac{\beta}{2}<y(\eta)<\beta\quad\forall\eta\in(0,\eta_\beta]. \tag{101}$$

An integration of (9) using (101), then gives,

$$\frac{\beta\eta}{2}<x(\eta)<\beta\eta\quad\forall\eta\in(0,\eta_\beta]. \tag{102}$$

It finally follows from (101) and (102), since $\eta_\beta \le \eta^*$, that,

$$x(\eta_\beta) < \beta\eta_\beta \le \beta\eta^* \le (1-p)^{\frac{1}{1-p}}, \quad y(\eta_\beta) > \frac{\beta}{2},$$

and so $(x(\eta_\beta), y(\eta_\beta)) \in \Omega$, which is a contradiction. We conclude that $\eta_\beta > \eta^*$, as required. □

Therefore, we have,

Corollary 12. *Let $(x,y) : [0,\eta^*] \to \mathbb{R}^2$ be a solution to (S) with zero-value $(0,\beta) \in \partial\Omega_1$ with η^* given by (98). Then,*

$$\frac{\beta\eta}{2} < x(\eta) < (1-p)^{1/(1-p)}, \quad \frac{\beta}{2} < y(\eta) < \beta \quad \forall \eta \in [0,\eta^*],$$

Proof. For cases (i) and (ii), the result follows from Proposition 11, with case (iii) following immediately. □

The a priori bounds in Corollary 12, allow us to establish the following local uniqueness result for (S) with zero-value $(0,\beta) \in \partial\Omega_1$. The proof is based on the uniqueness argument in [1].

Proposition 13. *The problem (S) with zero-value $(0,\beta) \in \partial\Omega_1$ has at most one solution on $[0,\eta^*]$, with $\eta^* > 0$ given by (98).*

Proof. To begin, fix $(0,\beta) \in \partial\Omega_1$. Suppose that $(x,y), (x^*,y^*) : [0,\eta^*] \to \mathbb{R}^2$ are solutions to (S) with zero-value $(0,\beta)$. It follows from Corollary 12 that

$$(x(\eta), y(\eta)), (x^*(\eta), y^*(\eta)) \in \bar{\Omega} \quad \forall \eta \in [0,\eta^*], \tag{103}$$

whilst from Corollary 12,

$$|x(\eta) - x^*(\eta)| < (1-p)^{1/(1-p)}, \quad |y(\eta) - y^*(\eta)| < \beta \quad \forall \eta \in [0,\eta^*]. \tag{104}$$

Additionally, we observe that for $(X,Y) \in [0,(1-p)^{1/(1-p)}] \times [0,\beta]$, then

$$X + X^p + Y < \left(2 + \beta^{1-p}\right)(X+Y)^p, \tag{105}$$

since $0 < p < 1$. Now, via (9) and (10) respectively, we have,

$$|x(\eta) - x^*(\eta)| \le \int_0^\eta |y(s) - y^*(s)| ds \tag{106}$$

$$|y(\eta) - y^*(\eta)| \le \int_0^\eta \left(\frac{1}{(1-p)} |x(s) - x^*(s)| + |x(s) - x^*(s)|^p + \frac{s}{2} |y(s) - y^*(s)| \right) ds \tag{107}$$

for all $\eta \in [0,\eta^*]$. We next introduce $v : [0,\eta^*] \to \mathbb{R}$ as,

$$v(\eta) = |x(\eta) - x^*(\eta)| + |y(\eta) - y^*(\eta)| \quad \forall \eta \in [0,\eta^*]. \tag{108}$$

Therefore, via (103)-(108), it follows that

$$\begin{aligned} v(\eta) &\leq \int_0^\eta \left(\frac{1}{(1-p)} |x(s) - x^*(s)| + |x(s) - x^*(s)|^p + \left(\frac{s}{2} + 1 \right) |y(s) - y^*(s)| \right) ds \\ &\leq \int_0^\eta \frac{1}{(1-p)} \left(\frac{\eta^*}{2} + 1 \right) \left(|x(s) - x^*(s)| + |x(s) - x^*(s)|^p + |y(s) - y^*(s)| \right) ds \\ &\leq \int_0^\eta \frac{1}{(1-p)} \left(\frac{\eta^*}{2} + 1 \right) (2 + \beta^{1-p}) \left(v(s) \right)^p ds \end{aligned} \tag{109}$$

for all $\eta \in [0, \eta^*]$, where the final inequality is due to (104) and (105). Also, via Corollary 12 and (98), η^* is dependent on p and β only, and hence, it follows from (109) that

$$v(\eta) \leq \int_0^\eta K(p, \beta) \left(v(s) \right)^p ds \tag{110}$$

for all $\eta \in [0, \eta^*]$, where the constant $K(p, \beta)$ is given by,

$$K(p, \beta) = \frac{1}{(1-p)} \left(\frac{\eta^*}{2} + 1 \right) (2 + \beta^{1-p}).$$

We now introduce the function $\bar{H} : [0, \eta^*] \to \overline{\mathbb{R}}_+$ given by,

$$\bar{H}(\eta) = \int_0^\eta K(p, \beta) \left(v(s) \right)^p ds \quad \forall \eta \in [0, \eta^*]. \tag{111}$$

It follows from (111) that $\bar{H}$ is non-negative, non-decreasing and differentiable on $[0, \eta^*]$, and via (110), satisfies

$$(\bar{H}(s))' \leq K(p, \beta) (\bar{H}(s))^p \quad \forall s \in [0, \eta^*]. \tag{112}$$

Upon integrating (112) from 0 to η, we obtain

$$\bar{H}(\eta) \leq \left((1-p) K(p, \beta) \eta \right)^{1/(1-p)} \quad \forall \eta \in [0, \eta^*] \tag{113}$$

and it follows from (113), (111) and (110) that

$$v(\eta) \leq \delta \quad \forall \eta \in [0, \eta_\delta], \tag{114}$$

where $\delta > 0$ is chosen sufficiently small so that

$$\eta_\delta = \frac{\delta^{1-p}}{(1-p) K(p, \beta)} < \eta^*.$$

Now, from Corollary 12, we have

$$\min\{x^*(\eta), x(\eta)\} \geq \frac{\beta \eta}{2} \quad \forall \eta \in [0, \eta^*]. \tag{115}$$

Moreover, it follows from (14), (115) and the mean value theorem, that there exists $\theta(s) \geq \min\{x^*(s), x(s)\}$, for which,

$$\begin{aligned}
&|Q_2(x(s), y(s), s) - Q_2(x^*(s), y^*(s), s)| \\
&\qquad \leq \frac{1}{(1-p)}|x(s) - x^*(s)| + |x(s)^p - x^*(s)^p| + \frac{s}{2}|y(s) - y^*(s)| \\
&\qquad \leq \frac{1}{(1-p)}|x(s) - x^*(s)| + p(\theta(s))^{p-1}|x(s) - x^*(s)| + \frac{\eta^*}{2}|y(s) - y^*(s)| \\
&\qquad \leq \left(\frac{1}{(1-p)} + p\left(\frac{\beta s}{2}\right)^{p-1}\right)|x(s) - x^*(s)| + \frac{\eta^*}{2}|y(s) - y^*(s)| \\
&\qquad \leq \left(\frac{1}{(1-p)} + p\left(\frac{\beta s}{2}\right)^{p-1} + \frac{\eta^*}{2}\right)v(s) \qquad (116)
\end{aligned}$$

for all $s \in (0, \eta^*]$. Now, via (9), (10), (14), (105), (116) and (114), we have,

$$\begin{aligned}
v(\eta) \leq &\int_0^{\eta} (|Q_1(x(s), y(s), s) - Q_1(x^*(s), y^*(s), s)| \\
&\qquad + |Q_2(x(s), y(s), s) - Q_2(x^*(s), y^*(s), s)|)\, ds \\
\leq &\int_0^{\eta_\delta} K(p, \beta)(v(s))^p ds + \int_{\eta_\delta}^{\eta} \left(1 + \frac{1}{(1-p)} + p\left(\frac{\beta s}{2}\right)^{p-1} + \frac{\eta^*}{2}\right)v(s) ds \\
\leq &\frac{\delta}{(1-p)} + \int_{\eta_\delta}^{\eta} \left(1 + \frac{1}{(1-p)} + p\left(\frac{\beta s}{2}\right)^{p-1} + \frac{\eta^*}{2}\right)v(s) ds \qquad (117)
\end{aligned}$$

for all $\eta \in [\eta_\delta, \eta^*]$. An application of Gronwall's Lemma [2, Corollary 6.2] to (117), gives

$$v(\eta) \leq \frac{\delta}{(1-p)} e^{\left(\eta^*\left(1 + \frac{1}{(1-p)} + \left(\frac{\beta\eta^*}{2}\right)^{p-1} + \frac{\eta^*}{2}\right)\right)} \qquad (118)$$

for all $\eta \in [\eta_\delta, \eta^*]$. Since v is non-negative and η^* is independent of δ, it follows from (118) and (114), upon letting $\delta \to 0$, that

$$v(\eta) = 0 \quad \forall \eta \in [0, \eta^*]. \qquad (119)$$

Finally, it follows from (119) and (108) that

$$(x(\eta), y(\eta)) = (x^*(\eta), y^*(\eta)) \quad \forall \eta \in [0, \eta^*],$$

as required. □

We can now state the following uniqueness result.

Lemma 14. *For each $(0, \beta) \in \partial\Omega_1$ then (S) with zero-value $(0, \beta)$ has exactly one solution $(x, y) : I \to \mathbb{R}^2$. This solution satisfies exactly one of the cases: (i) (with $I = [0, \eta_\beta + \epsilon_\beta]$), (ii) (with $I = [0, \eta_\beta + \epsilon_\beta]$) or (iii) (with $I = [0, \infty)$).*

Proof. We have established earlier that for each $(0,\beta) \in \partial\Omega_1$, then (S) with zero-value $(0,\beta)$ has at least one solution $(x,y): I \to \mathbb{R}^2$, and that the solution satisfies one of the cases (i)-(iii). It follows from Proposition 13 that this solution is unique for $\eta \in [0,\eta^*]$, (with η^* depending only upon β and p) and, moreover, in whichever case of (i)-(iii) it falls, that $(x(\eta),y(\eta)) \notin \{(0,\lambda) : \lambda \in \mathbb{R}\}$ for any $\eta \in I \setminus [0,\eta^*]$. Repeated application of the classical uniqueness theorem [6, Chapter 1, Theorem 2.2] then completes the uniqueness result for $\eta \in I \setminus [0,\eta^*]$. □

We immediately obtain a continuous dependence result for solutions of (S) with zero-value in $\partial\Omega_1$, namely,

Corollary 15. *Let $(0,\beta^*) \in \partial\Omega_1$ and suppose that the unique solution to (S) with zero-value $(0,\beta^*)$, say $(x^*,y^*): I \to \mathbb{R}$, satisfies case (i) or (ii), with $I = [0,\eta_{\beta^*}+\epsilon_{\beta^*}]$. Then, given $\epsilon' > 0$, there exists $\delta' > 0$ such that for all $\beta > 0$ satisfying $|\beta-\beta^*| < \delta'$, the corresponding unique solution to (S) with zero-value $(0,\beta)$, say $(x,y): I' \to \mathbb{R}$, has $I' = I$ and satisfies the corresponding case (i) or (ii), with,*

$$|x(\eta)-x^*(\eta)| + |y(\eta)-y^*(\eta)| < \epsilon' \quad \forall \eta \in I.$$

Proof. We first recall that (for a suitable choice of β^*) then

$$|x^*(\eta)| \le (1-p)^{\frac{1}{1-p}} + 1, \quad |y^*(\eta)| \le \beta^* + 1 \quad \forall \eta \in [0,\eta_{\beta^*}+\epsilon_{\beta^*}],$$

and, via (14), that $\mathbf{Q}(x,y,\eta)$ is continuous (and therefore bounded) on the rectangle

$$R = \left\{(x,y,\eta) : |x| \le (1-p)^{\frac{1}{1-p}} + 1, \quad |y| \le \beta^* + 1, \quad 0 \le \eta \le \eta_{\beta^*}+\epsilon_{\beta^*}\right\}.$$

The *uniqueness result* in Lemma 14 then allows for an application of the result [6, Theorem 4.3, pp. 59] which completes the proof. □

It is now convenient to introduce the three sets E_1, E_2 and E_3, where

$$E_1 = \{(0,\beta) \in \partial\Omega_1 \text{: the unique solution to (S)}$$
$$\text{with zero-value } (0,\beta) \text{ satisfies case (i)}\},$$

with E_2 and E_3 defined similarly for cases (ii) and (iii) respectively. It follows from Lemma 14 that

$$E_i \cap E_j = \varnothing \quad \text{for} \quad i,j = 1,2,3 \text{ with } i \ne j, \tag{120}$$

whilst

$$E_1 \cup E_2 \cup E_3 = \partial\Omega_1. \tag{121}$$

We now establish that E_1 and E_2 are both nonempty.

Proposition 16. *The set E_1 is non-empty and is such that $(0,\beta) \in E_1$ for each*

$$\beta > \sqrt{2\left(((1-p)^{1/(1-p)} - m_H)^2 - m_H^2\right)}, \tag{122}$$

with m_H given by (99).

Proof. Let $(x, y) : I \to \mathbb{R}^2$ be the unique solution to (S) with zero-value $(0, \beta) \in \partial\Omega_1$ and β satisfying (122). Since $(x(\eta), y(\eta)) \in \bar{\Omega}$ for all $\eta \in I'$ (where $I' = [0, \eta_\beta]$ for cases (i) and (ii), and $I' = [0, \infty)$ for case (iii)) then, via (9) and (10), we have,

$$\frac{\beta}{2} \leq y(\eta) \leq \beta, \quad x(\eta) \geq \frac{\beta\eta}{2} \quad \forall \eta \in [0, \bar{\eta}_\beta], \tag{123}$$

with,

$$\bar{\eta}_\beta = \begin{cases} \min\{\eta_\beta, \eta'_\beta\} & : \text{ cases (i) and (ii)} \\ \eta'_\beta & : \text{ case (iii)} \end{cases} \tag{124}$$

and

$$\eta'_\beta = \frac{2}{\beta}\left(m_H + \sqrt{m_H^2 + \frac{\beta^2}{2}}\right).$$

Now suppose case (iii) occurs, then $(x(\eta'_\beta), y(\eta'_\beta)) \in \Omega$. However,

$$x(\eta'_\beta) \geq \frac{\beta\eta'_\beta}{2} = m_H + \sqrt{m_H^2 + \frac{\beta^2}{2}} > (1-p)^{\frac{1}{1-p}}$$

via (123) and (122), and we arrive at a contradiction. We can therefore eliminate case (iii). Next suppose case (ii) occurs. It follows from $(123)_2$ and (124) that $\eta_\beta \leq \eta'_\beta$, and so $\bar{\eta}_\beta = \eta_\beta$. Thus, via $(123)_1$,

$$y(\eta_\beta) \geq \frac{\beta}{2} > 0.$$

However, in case (ii), $y(\eta_\beta) = 0$, and we arrive at a contradiction. We conclude finally that case (i) must occur, as required. □

We can also establish a similar result for E_2.

Proposition 17. *The set E_2 is non-empty and is such that $(0, \beta) \in E_2$ for each*

$$0 < \beta < \sqrt{\frac{(1-p)^{2/(1-p)}}{(1+p)}}. \tag{125}$$

Proof. It follows from (16)-(21) that for β satisfying the inequality (125), then $(0, \beta) \in D_{c^*(p)}$. It then follows from Corollary 4 that (S) with zero-value $(0, \beta)$ has a global solution which lies in D_{c_β} for all $\eta \in (0, \infty)$ with $c_\beta = V(0, \beta) < c^*(p)$, and so the solution to (S) in $\eta \geq 0$ must satisfy case (ii). Therefore, $(0, \beta) \in E_2$, as required. □

We next establish that both E_1 and E_2 are open subsets of $\partial\Omega_1$.

Proposition 18. *The sets E_1 and E_2 are open subsets of $\partial\Omega_1$.*

Proof. We will prove the result for E_1. The proof for E_2 is similar. Let $(0, \beta^*) \in E_1$. Then, via Lemma 14, (S) with zero-value $(0, \beta^*)$ has a unique solution $(x^*, y^*) : [0, \eta_{\beta^*} + \epsilon_{\beta^*}] \to \mathbb{R}^2$, with

$$(x^*(\eta), y^*(\eta)) \in \Omega \quad \forall \eta \in (0, \eta_{\beta^*}) \tag{126}$$

and

$$(x^*(\eta_{\beta^*}), y^*(\eta_{\beta^*})) = ((1-p)^{1/(1-p)}, y_{\beta^*}) \tag{127}$$

for some $0 < y_{\beta^*} < \beta^*$, whilst

$$(x^*(\eta), y^*(\eta)) \notin \bar{\Omega} \quad \forall \eta \in (\eta_{\beta^*}, \eta_{\beta^*} + \epsilon_{\beta^*}]. \tag{128}$$

Now consider the family of open balls

$$B(x^*(\eta), y^*(\eta); \epsilon') \text{ with } \eta \in [0, \eta_{\beta^*} + \epsilon_{\beta^*}]$$

and via (126)-(128), choose ϵ' sufficiently small so that

$$B(x^*(\eta_{\beta^*} + \epsilon_{\beta^*}), y^*(\eta_{\beta^*} + \epsilon_{\beta^*}); \epsilon') \cap \bar{\Omega} = \varnothing \tag{129}$$

and

$$\bigcup_{\lambda \in [0, \eta_{\beta^*} + \epsilon_{\beta^*}]} B(x^*(\lambda), y^*(\lambda); \epsilon') \cap (\partial\Omega \setminus \partial\Omega_1) \subset \{((1-p)^{1/(1-p)}, \lambda) : \lambda > 0\}. \tag{130}$$

It then follows from Corollary 15 that there exists $\delta' > 0$ such that the corresponding unique solution to (S) with zero-value $(0, \beta) \in \partial\Omega_1$, satisfying $|\beta - \beta^*| < \delta'$, say $(x, y) : [0, \eta_{\beta^*} + \epsilon_{\beta^*}] \to \mathbb{R}^2$ has

$$(x(\eta), y(\eta)) \in \bigcup_{\lambda \in [0, \eta_{\beta^*} + \epsilon_{\beta^*}]} B(x^*(\lambda), y^*(\lambda); \epsilon') \quad \forall \eta \in [0, \eta_{\beta^*} + \epsilon_{\beta^*}] \tag{131}$$

Therefore, via (129)-(131), $\{(0, \beta) : |\beta - \beta^*| < \delta'\} \subseteq E_1$, and so E_1 is an open subset of $\partial\Omega_1$, as required. □

Finally, we have

Corollary 19. *The set E_3 is a non-empty closed subset of $\partial\Omega_1$.*

Proof. Via Propositions 16 and 17, E_1 and E_2 are both nonempty subsets of $\partial\Omega_1$. Moreover, via (120) E_1 and E_2 are disjoint. Suppose that E_3 is empty, then via (121) and Proposition 18, E_1 and E_2 form an open partition of $\partial\Omega_1$. However, $\partial\Omega_1$ is a connected subset of $\mathbb{R}^2$, and we arrive at a contradiction. Hence E_3 must be nonempty. Finally, $E_3 = \partial\Omega_1 \setminus (E_1 \cup E_2)$ and is therefore a closed subset of $\partial\Omega_1$. □

Remark 3. *In Corollary 19, the existence of at least one point in E_3 has been established. However, it has not been established that this is the only point in E_3.*

To conclude this section, we arrive at our main result, namely,

Theorem 20. *There exists a solution* $(x,y):\mathbb{R}\to\mathbb{R}^2$ *to (S) with zero-value* $(0,\beta)\in\partial\Omega_1$*, for some*

$$\sqrt{\frac{(1-p)^{2/(1-p)}}{(1+p)}}\le\beta\le\sqrt{2\left(((1-p)^{1/(1-p)}-m_H)^2-m_H^2\right)},$$

which satisfies

$$(x(\eta),y(\eta))\to(\pm(1-p)^{1/(1-p)},0)\ \text{ as }\eta\to\pm\infty \tag{132}$$

and

$$|x(\eta)|<(1-p)^{1/(1-p)},\quad 0<y(\eta)\le\beta\quad\forall\eta\in\mathbb{R}. \tag{133}$$

Proof. It follows directly from Corollary 19 and (iii) that there exists $(x^*,y^*):[0,\infty)\to\mathbb{R}^2$ which is a solution to (S) with zero-value $(0,\beta^*)$, such that

$$(x^*(\eta),y^*(\eta))\to((1-p)^{1/(1-p)},0)\text{ as }\eta\to\infty, \tag{134}$$

$$(x^*(\eta),y^*(\eta))\in\Omega\quad\forall\eta\in(0,\infty). \tag{135}$$

It follows from (125) and (122), that

$$\sqrt{\frac{(1-p)^{1/(1-p)}}{(1+p)}}\le\beta^*\le\sqrt{2\left(((1-p)^{1/(1-p)}-m_H)^2-m_H^2\right)}.$$

Now, define the function $(x,y):\mathbb{R}\to\mathbb{R}^2$ to be

$$(x(\eta),y(\eta))=\begin{cases}(x^*(\eta),y^*(\eta)) & ;\eta\in[0,\infty)\\ (-x^*(-\eta),y^*(-\eta)) & ;\eta\in(-\infty,0).\end{cases} \tag{136}$$

It follows from (136) that $(x,y):\mathbb{R}\to\mathbb{R}^2$ is a solution to (S) with zero-value $(0,\beta^*)$, and via (94) and (iii), (since $y(\eta)$ is monotone decreasing for $\eta\in(0,\infty)$) that this solution satisfies (132) and (133). □

We conclude from Theorem 20 that the problem (S) has at least one heteroclinic connection from the equilibrium point $(-(1-p)^{1/(1-p)},0)$ $(\eta=-\infty)$ to the equilibrium point $((1-p)^{1/(1-p)},0)$ $(\eta=\infty)$, which we denote by $w_{\beta^*}:\mathbb{R}\to\mathbb{R}$. Here $w=w_{\beta^*}(\eta)$, $\eta\in\mathbb{R}$, has zero-value $w(0)=0$, $w'(0)=\beta^*$ for some

$$\sqrt{\frac{(1-p)^{2/(1-p)}}{(1+p)}}\le\beta^*\le\sqrt{2\left(((1-p)^{1/(1-p)}-m_H)^2-m_H^2\right)},$$

and

$$|w_{\beta^*}(\eta)|<(1-p)^{1/(1-p)},\quad 0<w'_{\beta^*}(\eta)\le\beta^*\quad\forall\eta\in\mathbb{R},$$

recalling also, that $w_{\beta^*}(\eta)$ is an odd function of $\eta\in\mathbb{R}$. Finally, a straightforward linearization as $|\eta|\to\infty$ establishes that,

$$w_{\beta^*}(\eta)\sim\pm(1-p)^{1/(1-p)}-\frac{A_\infty}{\eta^3}e^{-\frac{1}{4}\eta^2}\quad\text{as}\quad\eta\to\pm\infty,$$

with A_∞ being a globally determined constant.

4.2 Front Solutions to [CP]

Following Theorem 20, with $\beta = \beta^*$ we have constructed the front-like global solution $u_{\beta^*} : \mathbb{R} \times [0, \infty) \to \mathbb{R}$ to [CP], namely,

$$u_{\beta^*}(x,t) = \begin{cases} t^{\frac{1}{(1-p)}} w_{\beta^*}\left(\frac{x}{t^{1/2}}\right) & , (x,t) \in \mathbb{R} \times (0,\infty) \\ 0 & , (x,t) \in \mathbb{R} \times \{0\}. \end{cases} \tag{137}$$

We again observe that $u_{\beta^*}(x - x_0, t)$ is also a global solution to [CP] for any fixed $x_0 \in \mathbb{R}$. In addition, following Section 3.3, we conclude that, for any $\tau > 0$, $u^\tau_{\beta^*} : \mathbb{R} \times [0, \infty) \to \mathbb{R}$ such that

$$u^\tau_{\beta^*}(x,t) = \begin{cases} (t-\tau)^{\frac{1}{(1-p)}} w_{\beta^*}\left(\frac{x}{(t-\tau)^{1/2}}\right) & , (x,t) \in \mathbb{R} \times (\tau,\infty) \\ 0 & , (x,t) \in \mathbb{R} \times [0,\tau] \end{cases}$$

is also a front-like global solution to [CP].

5 Discussion

There are two questions that arise naturally from this study. The first being how one can rigorously establish the decay rate of the homoclinic solutions $w : \mathbb{R} \to \mathbb{R}$ to (S) as $\eta \to \pm\infty$, that is suggested by (87) and (88); the second being whether or not for the problem (S), there is a unique heteroclinic connection from the equilibrium point $(-(1-p)^{1/(1-p)}, 0)$ to the equilibrium point $((1-p)^{1/(1-p)}, 0)$ which has zero value in $\partial\Omega_1$ (Theorem 20 guarantees that there exists at least one connection).

References

[1] J. Aguirre and M. Escobedo, "A Cauchy problem for $u_t - \triangle u = u^p$ with $0 < p < 1$. Asymptotic behavior of solutions." Annales Faculté des Sciences de Toulouse, **8**, 2, (1986), 175-203.

[2] H. Amann, *Ordinary differential equations: an introduction to nonlinear analysis.* (de Gruyter, Berlin, 1990).

[3] J. Ball, "Remarks on blow-up and nonexistence theorems for nonlinear evolution equations." Quart. J. Math. Oxford Ser. (2), **28**, 112, (1977), 473-486.

[4] T. Cazenave, F. Dickstein and F. B. Weissler, "Sign-changing stationary solutions and blowup for the nonlinear heat equation in a ball." Math. Ann., **344**, 2, (2009), 431-449.

[5] T. Cazenave, F. Dickstein and F. B. Weissler, "Multi-scale multi-profile global solutions of parabolic equations in $\mathbb{R}^N$." Discrete Contin. Dyn. Syst. Ser. **5**, 3, (2012), 449-472.

[6] E. A. Coddington and N. Levinson, *Theory of Ordinary Differential Equations.* (McGraw-Hill, 1955, London).

[7] K. Deng and H. A. Levine, "The role of critrical exponents in blow-up theorem:the sequel." J. Math. Anal. Appl. **243**, no. 1, (2000), 85-126.

[8] C. Dohmen and M. Hirose, "Structure of positive radial solutions to the haraux weissler equation." Nonlinear Anal. **33**, no. 1, (1998), 51-69.

[9] M. Escobedo and O. Kavian, "Variational problems related to self-similar solutions of the heat equation." Nonlinear Anal. **11**, no. 10, (1987), 1103-1133.

[10] A. Haraux and F. B. Weissler, "Non-uniqueness for a Semilinear Initial Value Problem." Indiana Uni. Math. J., **31**, 2, (1982), 167-189.

[11] M. Hirose and E. Yanagida, "Global Structure of Self-Similar Solutions in a Semilinear Parabolic Equation." J. Math. Anal. Appl. **244**, no. 2, (2000), 348368.

[12] A. C. King and D. J. Needham, "On a singular initial-boundary value problem for a reaction-diffusion equation arising from a simple model of isothermal chemical autocatalysis. "Proc. R. Soc. Lond, A" **437**, 1901, (1992) 657-671.

[13] H. A. Levine, "The role of critical exponents in blowup theorems." SIAM Rev. **342**, no. 2, (1990) 262-288.

[14] P. M. McCabe , J. A. Leach and D. J. Needham, "A note on the non-existence of permanent form travelling wave solutions in a class of singular reaction-diffusion problems.", Dyn. Syst., **17**, 2, (2002), 131-135.

[15] J. C. Meyer and D. J. Needham, "Extended weak maximum principles for parabolic partial differential inequalities on unbounded domains." Proc. R. Soc. Lond. A **470**, no. 2167, (2014).

[16] J. C. Meyer and D. J. Needham, *The Cauchy Problem for Non-Lipschitz Semi-Linear Parabolic Partial Differential Equations.* LMS Lecture Notes in Mathematics Series: (Cambridge University Press, Cambridge , 2015).

[17] P. D. Miller, *Applied Asymptotic Analysis.* (AMS, 2006, Rhode Island).

[18] N. Mizoguchi and E. Yanagida, "Critical exponents for the blow-up of solutions with sign changes in a semilinear parabolic equation" Math. Ann., **307**, (1997), 663-675.

[19] N. Mizoguchi and E. Yanagida, "Critical exponents for the blow-up of solutions with sign changes in a semilinear parabolic equation, II" J. Differential. Equations 145, no. 2, (1998), 295-331.

[20] D. J. Needham, "On the global existence of solutions to a singular semilinear parabolic equation arising from the study of autocatalytic chemical kinetics." ZAMP **43**, 3, (1992), 471-480.

[21] D. J. Needham and P. G. Chamberlain, "Global similarity solutions to a class of semilinear parabolic equations: existence, bifurcations and asymptotics." Proc. R. Soc. Lond. A. **454**, no. 1975, (1998), 1933-1959.

[22] P. Poláčik, "Symmetry properties of positive solutions of parabolic equations: a survey". In *Recent progress on reaction-diffusion systems and viscosity solutions* (ed. E.Y. Du, H. Ishii and W-Y. Lin), Singapore: World Sci. Pub. Co. 170208, World Sci. Publ., (2009), Hackensack, NJ.

[23] P. Poláčik, "Symmetry properties of positive solutions of parabolic equations on $\mathbb{R}^N$: I. Assymptotic symmetry for the Cauchy problem". Comm. Partial Differential Equations 30 (2005), no. 10-12, 1567-1593.

[24] P. Poláčik and E. Yanagida, "On bounded and unbounded global solutions of a supercritical semi-linear heat equation." Math. Ann. **327**, (2003) 745-771.

[25] N. Shioji and K. Watanabe, "A generalised Pohozaev identity and uniqueness of positive radial solutions of $\triangle u + g(r)u + h(r)u^p = 0$." J. Differential Equations **255** (2013), 4448-4475.

[26] F. B. Weissler, "Existence and nonexistence of global solutions for a semilinear heat equation." Israel J. Math., **38**, (1981), 29-40.

[27] F. B. Weissler, "Local existence and nonexistence for semilinear parabolic equations in L^p." Indiana Univ. Math. J. **29**, (1980), 79-102.

J. C. Meyer, School of Mathematics, Watson Building, University of Birmingham, Birmingham, UK, B15 2TT
E-mail address: J.C.Meyer@bham.ac.uk

D. J. Needham, School of Mathematics, Watson Building, University of Birmingham, Birmingham, UK, B15 2TT
E-mail address: D.J.Needham@bham.ac.uk